



LESCO®

#279974



DIMENSION™ CRABGRASS PRE-EMERGENT

PLUS FERTILIZER **19-0-7**
Pasto de cuaresma de dimensión
pre-emergente más abono



Made in USA

PLUS FERTILIZER
Pasto de cuaresma de dimensión
pre-emergente más abono

DIMENSION™
CRABGRASS
PRE-EMERGENT **19-0-7**



Made in USA

PLUS FERTILIZER
Pasto de cuaresma de dimensión
pre-emergente más abono



9


#279974

19-0-7

N **NITROGEN**
For vigorous blade, leaf and stem growth and dark green color
NITRÓGENO
Para que el tallo y hojas tengan un crecimiento vigoroso y un color verde oscuro

P **PHOSPHATE**
Contains no phosphorus
FÓSFORO
Contiene ningún fósforo

K **POTASH**
Promotes strong roots and improves turf hardness
POTASA
Promueve raíces fuertes y mejora el vigor del césped.



Controls yellow foxtail, buttercup oxalis, and spotted spurge
Controla foxtail amarillo, oxalis buttercup y spurge manchado.




Nationwide
En todo el país.



Provides up to 4 months of weed control if applied according to label directions
Proporciona hasta 4 meses de control de pasto de cuaresma y otras hierbas de césped y de hoja ancha.



Spring
Primavera



Summer
Verano



Fall
Otoño

DIMENSION™
CRABGRASS
PRE-EMERGENT **19-0-7**



Made in USA



Covers up to
13,000
Sq. Ft.

ACTIVE INGREDIENTS
Dithiopyr: S,S'-dimethyl 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-3,5-pyridinedicarbothioate.....0.15%
OTHER INGREDIENTS.....99.85%
TOTAL.....100.00%
Product protected by U.S. Patent No. 4,692,184. Other patents pending.
KEEP OUT OF REACH OF CHILDREN
CAUTION | PRECAUCION
If you do not understand this label, find someone to explain it to you in detail.
Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS
Read the entire label before using this product.
Use only according to label instructions.

#080311 | NET. WT. 50 LB (22.7 KG)



DIMENSION™
CRABGRASS
PRE-EMERGENT **19-0-7**



Made in USA

PLUS FERTILIZER
Pasto de cuaresma de dimensión
pre-emergente más abono



DIMENSION™
CRABGRASS 
PRE-EMERGENT
PLUS FERTILIZER

19-0-7

LESCO® Crabgrass Pre-Emergent Plus Fertilizer is a premium quality combination product that provides both weed control and the required nutrients for optimum turfgrass growth all in a single product. It contains Dimension herbicide, a selective herbicide that can be applied up to 4 weeks later than other crabgrass preventers and provides up to 4 months control of crabgrass and other grasses and broadleaf weeds when applied prior to their germination in lawns and ornamental turfgrasses.

GUARANTEED ANALYSIS	
Total Nitrogen (N).....	19.00%
19.00% Urea Nitrogen*	
Soluble Potash (K ₂ O).....	7.00%
Derived From: Muriate of Potash, Polymer Coated Urea, Urea	
*5.13% Slowly Available Nitrogen from Polymer Coated Urea	F1560

SUGGESTED ROTARY SPREADER SETTINGS	
ROTARY SPREADER Recommended Rates in lb. per 1,000 sq. ft.	3.82 lb.
LESCO® Commercial Calibration Gauge (80 pound hopper and larger)	#15
Vigoro® 4300 Walk-Behind Rotary Spreader	# 4
Scotts® Broadcast/Rotary Spreaders (Scotts® Basic®, Standard™, Deluxe EdgeGuard® or SpeedyGreen®)	# 6

AN EXPANDED SPREADER SETTING LISTING CAN BE FOUND AT: www.LESCO.com

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION:

Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

ENVIRONMENTAL HAZARDS:

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment wash waters.

KEEP OUT OF REACH OF CHILDREN
CAUTION/PRECAUCIÓN
 If you do not understand this label, find someone to explain it to you in detail. Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

EPA Reg. No. 10404-86 | EPA Est. No. 59144-GA-1(A)
 EPA Est. No.32802-MO-1(B) | EPA Est. No. 961-PA-01(L)
 EPA Est. No.961-IL-01(D) | EPA Est. No. 37351-CO-1 (E)
 EPA Est. No. 8378-IN-1(K) | EPA Est. No. 35497-OR-1(H)
 EPA Est. No.49991-TX-2(Q) | EPA Est. No. 82757-OH-001 (M)
 EPA Est. No. 82757-FL-001 (S) | EPA Est. No.82757-MA-002 (H)
 First letter in lot code indicates manufacturing site.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. **Pesticide Storage:** Store this product only in its original container in a dry, cool, secured storage area. **Pesticide Disposal:** Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal State or local procedures. Or call [1-800-CLEANUP] for disposal instructions. Never place unused product down any indoor or outdoor drain. **Container Disposal:** Non-refillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. **Spill:** In case of spill, sweep up material and dispose of material according to "pesticide disposal" directions listed above.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. The directions for use of this product should be followed carefully. It is impossible to eliminate weather, presence of other materials or other influencing factors, which are beyond the control of LESCO, Inc. All such risks shall be assumed by Buyer, and Buyer agrees to hold LESCO, Inc. harmless for any claims relating to such factors. LESCO, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the directions of use, subject to the inherent risks referred to above, when used in accordance with instruction, or under abnormal conditions or under OR OF FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE. In no event shall LESCO, Inc. be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER AND THE EXCLUSIVE LIABILITY OF LESCO, INC. FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY INC. REPLACEMENT OF THE PRODUCT. LESCO, Inc. offers this product, and Buyer accepts it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability; this product is available on the Internet at <http://www.aapfco.org/metals.htm>. Information concerning the raw material composing this product can be obtained by writing to: LESCO, Inc Attn: RA Dept 1385 East 36th Street, Cleveland, Ohio 44114-4114, referring to the item number found on this bag. | LESCO and Poly Plus are registered trademarks and the sweeping design is a trademark of LESCO® Technologies, LLC. SCOTTS®, SpeedyGreen® And EasyGreen® are registered trademarks of SCOTT® Company; Vigoro® is a registered trademark of the United Industries Corporation. DIMENSION is a registered trademark of Dow AgroSciences LLC.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner that is inconsistent with its labeling. **Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production or for research purposes.**

GENERAL INFORMATION - Apply to well established lawns and ornamental turfgrasses listed below. (See "USE ON THE FOLLOWING TURFGRASSES" section for a list of the tolerant turfgrasses). This product will not harm nearby established ornamentals when used according to directions given below.

USE PRECAUTIONS - This product should only be applied to lawns and ornamental turfgrasses that are well established. Use of this product on lawns and ornamental turfgrasses that are not well established, or have been weakened by stress due to mechanical operations, pests, excessive temperatures, or lack of moisture, may increase the chances for turfgrass injury. Do not apply to lawns or turfgrasses that are under stress.

USE ON THE FOLLOWING TURFGRASSES - This product can be applied to the following turfgrasses when used as directed:

Cool-Season Turfgrasses:
 Creeping Bentgrass | Fine Fescue | Kentucky Bluegrass | Perennial Ryegrass | Tall Fescue

Warm-Season Turfgrasses:
 Bahiagrass | Bermudagrass | Buffalograss | Carpetgrass | Centipedegrass | St. Augustinegrass | Zoysiagrass

This product should be applied in the spring or fall prior to weed seed germination in established and newly germinated lawns and ornamental turfgrasses. It can be applied up to 4 weeks later than other crabgrass preventers and still provide season - long weed control. This is due to the dithiopyr herbicide component of this combination product. This product provides both preventative (pre-emergence) control of crabgrass and other weeds then applied prior to their germination, and post-emergence control of seedling crabgrass. Application of this product later than 4 weeks after crabgrass germination will result in reduced control of crabgrass. The post-emergence control provided by this product is limited to crabgrass only and will provide control of crabgrass when applied up to 4 weeks after crabgrass germination. Application of this product later than 4 weeks after crabgrass germination will result in reduced control of crabgrass. For control of other grass and broadleaf weeds this product must be applied as a preventative treatment before their germination. Only crabgrass will be controlled with delayed applications after germination. Tips for improving weed control: For best results apply this product within few days after mowing and delay further mowing after the application until the lawn or ornamental turfgrass has been watered or received significant rainfall to dissolve the fertilizer granules. Improved weed control may be observed when treated lawn or ornamental turfgrass is watered or receives rainfall with a few days after application of this product.

RESEEDING, OVERSEEDING OR SPRIGGING - Reseeding, overseeding, or sprigging of treated areas with this product should be delayed until 12 weeks from the time of application. Reseeding, overseeding or sprigging before 12 weeks after application may prevent establishment of desirable turfgrasses. When reseeding or overseeding, proper cultural practice such as soil cultivation, irrigation and fertilization should be followed. However, if overseeding with perennial ryegrass in the fall, overseed 8 weeks or later after a single application. For best results, use mechanical or power seeding equipment (silt seeders) designed to give good seed to soil contact. To establish a new lawn of Kentucky Bluegrass, Perennial Ryegrass and/or Tall Fescue, this product may be applied 2 weeks after the first sign of germination. For other tolerant turfgrasses listed on this label, wait until the grass is well established (2 mowing) before applying this product.

- DO NOT apply to flowers, vegetables or fruit trees.
- DO NOT use clippings from treated turf for mulching around flowers, vegetables or fruit trees.
- DO NOT apply this product to pastures.
- Keep people and pets off treated areas until dust has settled.
- DO NOT apply this product within 8 weeks of seeding.
- DO NOT apply more than 3.82 pounds of this product per 1,000 sq. ft. per application.
- In the State of New York, make only one application per year. In other states, do not make more than 3 applications per year spaced at least 2-4 months apart according to "Region."

WEEDS CONTROLLED - This product will provide control of crabgrass and other grasses and broadleaf weeds listed below when applied prior to germination in established lawns and ornamental turfgrasses.

Grass Weeds
 Annual Bluegrass | Annual Ryegrass | Barnyardgrass | Crabgrass | Crowfootgrass | Goosegrass | Green Foxtail | Kikuyugrass (seedling) | Perennial Ryegrass | Smutgrass (seedling) | Yellow Foxtail

Broadleaf Weeds
 Bittercress | Buttercup Oxalis | Carolina Geranium | Chickweed | Common Lespedeza | Common Purslane | Corn Speedwell
 Creeping Oxalis | Henbit | Parsley-piert | Pineappleweed | Prostrate Spurge | Spotted Spurge | Yellow Woodsorrel

WHEN TO APPLY - Application Timings - This product may be applied either in the spring or in the fall for the control of weeds. When this product is applied in the fall, a sequential or second application should be made the following spring to provide season-long control of weeds. If this product is applied in the fall, DO NOT reseed, overseed or sprig turfgrasses before 8 weeks after application of this product due to potential turfgrass injury to the newly reseeded, overseeded or sprigged turfgrasses.

LENGTH OF WEED CONTROL - This product may provide up to 4 months of weed control if applied according to label directions. The actual length of control will vary depending on use rate, weather conditions, condition of the turfgrass, intensity of weed pressure, and the location or "Region" within the United States where this product is being applied. Please check the map below to determine which "region" of the U.S. that applies to your particular use area, and find the number of pounds of this product needed in your area and on the length of weed controlled given in the Recommended Use Rates For This Product.

POSTEMERGENCE CRABGRASS CONTROL - For post-emergent crabgrass control, apply this product at the recommended rates when crabgrass is not beyond 3-4 leaf stage. Applications will control existing crabgrass up to the 4-leaf stage and prevent further growth.


HOW TO APPLY - APPLICATION EQUIPMENT AND INSTRUCTIONS

This product can be applied with a rotary or drop type spreader designed to apply granular fertilizers. For best results, apply the product evenly and uniformly over the turfgrass area avoiding streaking, skips or overlaps. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions. Adjust the spreader setting such that the spreader delivers the appropriate application rates of this product as recommended below.

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 sq. ft. per growing season. When used as directed, this product will supplement your regular fertilizer program by providing 0.73 lb of nitrogen per 1,000 sq. ft. at the 3.82 lb. rate and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program. *Based on Inert Fertilizer claimed.

A more uniform applications be made by spreading one-half of this product at a right angle to the previous application direction. Check application equipment frequently to ensure equipment is functioning properly and applying a uniform distribution of product. *0.5 pounds of nitrogen per 1,000 sq. ft. at the 2.67 lb. rate.

NOTICE: Before using this product, read the entire Precautionary statements, Directions for Use, Storage and Disposal Instructions and Warranty Statements. If the Warranty Statements are not acceptable, return the product unopened within 30 days of purchase to the place of purchase.



RECOMMENDED USE RATES FOR THIS PRODUCT LENGTH OF WEED CONTROL		
Region of U.S.	3-4 Months Control	Early Post Emergence Control
North	2.67 (0.174 lb. a/acre)*	3.82 (0.25lb.a/acre)*
South/West	3.82 (0.25lb. a/acre)*	3.82 (0.25lb.a/acre)*
(Pounds of this product to apply per 1,000 sq ft of turfgrass) *Use the higher rate in areas of high weed pressure.		

#080311

NET. WT. 50 LB (22.7 KG)

Covers up to
13,000
 Sq. Ft.

FOR PRODUCT ASSISTANCE, CALL LESCO
1-800-321-5325
 OR VISIT US AT WWW.LESCO.COM

FREIGHT SHIPPING BAG
 Meeting Requirements of
 APPLICABLE FREIGHT CLASSIFICATION
 Guaranteed by
 TRINITY PACKAGING CORPORATION

F1560 | Rev. Fall 2015
 Manufactured for LESCO, Inc.
 1385 East 36th St.
 Cleveland, OH 44114-4114

18 X 29